Ablumir
Built Around Immunology

Anti-Cdc2 Rabbit mAb [13A11]

Catalog No.: A52072

Basic Information

Catalog No. (Cat.No.)	A52072
Host species	Rabbit
Reactivity	Human,Mouse
Applications	WB,IHC,IP,IF,ELISA
Synonyms	Cdc2,Cell division cycle protein 2 homolog (S. cerevisiae),p34cdc2,MPM2 antigen

Product Information

Target	Cdc2
Molecular Weight(Calculated)	34 kDa
Molecular Weight(Observed)	34 kDa
Isotype	IgG,Kappa
Conjugation	NA
Concentration	0.2 mg/ml
Clonality	Monoclonal
Clone No.	13A11
RRID	NA
Storage buffer	50% Glycerol, 0.05% BSA and 0.05% preservative in PBS (pH 7.2-7.4)
Storage	Stored at -20°C for 1 year. Avoid repeated freeze / thaw cycles.
Shipped at conditions	Frozen
Form	Liquid

Ablumir
Built Around Immunology

Product Usage Information

Application	Dilution
WB	1:2000-1:10000
IHC	1:1000-1:4000
IP	1:50-1:200
IF	1:200-1:1000
ELISA	1:5000-1:20000

Target Information

Protein Name	Cyclin-dependent kinase 1
Gene Name	*CDK1*
Domain	NA
Background	Cdc2 (Cell-division-cycle 2, CDK1) belongs to cyclin-dependent kinase family and serves as master regulator for G2/M transition and mitosis onset. Cdc2 associates with Cyclin B to form maturation-promoting factor (MPF). Its kinase activity is tightly governed by layered phosphorylation. Wee1 introduces inhibitory phosphorylation to keep Cdc2 silent; Cdc25C removes these inhibitory phosphates to activate MPF, triggering nuclear envelope breakdown, spindle assembly and other mitotic events. Cyclin B degradation after mitosis terminates Cdc2 activity. Dysregulated Cdc2 activation causes cell-cycle aberration and genomic instability, and its up-regulation is detected in various cancers.