Ablumir
Built Around Immunology

Anti-p38 MAPK (phospho Thr180/Tyr182) Rabbit mAb [58F80]

Catalog No.: A52257

Basic Information

Catalog No. (Cat.No.)	A52257
Host species	Rabbit
Reactivity	Human,Mouse,Rat
Applications	WB,IHC,IP,IF,ELISA
Synonyms	p38 MAPK (phospho Thr180/Tyr182),p-p38(Thr180/Tyr182),p38α MAPK,p38-alpha

Product Information

Target	p38 MAPK (phospho Thr180/Tyr182)
Molecular Weight(Calculated)	44 kDa
Molecular Weight(Observed)	44 kDa
Isotype	IgG,Kappa
Conjugation	NA
Concentration	Please check the information on the tube.
Clonality	Monoclonal
Clone No.	58F80
RRID	NA
Storage buffer	50% Glycerol, 0.05% BSA and 0.05% preservative in PBS (pH 7.2-7.4)
Storage	Stored at -20°C for 1 year. Avoid repeated freeze / thaw cycles.
Shipped at conditions	Frozen
Form	Liquid

Ablumir
Built Around Immunology

Product Usage Information

Application	Dilution
WB	1:2000-1:10000
IHC	1:200-1:1000
IP	1:50-1:200
IF	1:200-1:1000
ELISA	1:5000-1:20000

Target Information

Protein Name	p38 MAPK (phospho Thr180/Tyr182)
Gene Name	*MAPK14*
Domain	NA
Background	p38 MAPK (Mitogen-activated protein kinase 14) belongs to MAP kinase family and transduces cellular stress, inflammatory, apoptotic and differentiation signals. p38 is a serine-threonine kinase activated via upstream MAP2K-mediated dual phosphorylation upon pro-inflammatory cytokines, oxidative stress, UV irradiation and osmotic challenge. Activated p38 translocates into nucleus and phosphorylates transcription factors to modulate transcription of pro-inflammatory, apoptotic and cell-cycle target genes. It mediates innate immunity and cytokine production, as well as cell differentiation and senescence. Sustained p38 signaling contributes to autoimmunity, inflammatory disorders and tumor progression.