Ablumir
Built Around Immunology

Anti-AMPKα1/2 (phospho Thr183/Thr172) Rabbit pAb (A72007)

Catalog No.: A72007

Basic Information

Catalog No. (Cat.No.)	A72007
Host species	Rabbit
Reactivity	Human,Mouse,Rat,Monkey,Pig,Marsupenaeus japonicus
Applications	WB,IHC,IF,ELISA
Synonyms	AMPKα1/2 (phospho Thr183/Thr172),p-AMPK-α1/2(Thr183/172),p-AMPKα(T183/T172),Phospho-AMPKα

Product Information

Target	AMPKα1/2 (phospho Thr183/Thr172)
Molecular Weight(Calculated)	62 kDa
Molecular Weight(Observed)	63 kDa
Isotype	IgG
Conjugation	NA
Concentration	1 mg/ml
Clonality	Polyclonal
Clone No.	NA
RRID	NA
Storage buffer	50% Glycerol, 0.5% BSA and 0.02% preservative in PBS (pH 7.2-7.4)
Storage	Stored at -20°C for 1 year. Avoid repeated freeze / thaw cycles.
Shipped at conditions	Frozen
Form	Liquid

Ablumir
Built Around Immunology

Product Usage Information

Application	Dilution
WB	1:500-1:2000
IHC	1:100-1:300
IF	1:50-1:200
ELISA	1:40000

Target Information

Protein Name	AMPKα1/2 (phospho Thr183/Thr172)
Gene Name	*PRKAA1 / PRKAA2*
Domain	human AMPK alpha,around phosphorylation site of Thr172, aa 140-189
Background	Phospho-AMPKα (Thr183/Thr172) corresponds to the conserved activating phosphorylation sites of AMPKα1 and AMPKα2 subunits, which are the hallmark modifications for full AMPK activation. Thr172 on AMPKα2 and Thr183 on AMPKα1 are located in the kinase activation loop, and their phosphorylation is a prerequisite for initiating AMPK catalytic activity and downstream signaling. As a central energy sensor, activated AMPK suppresses anabolic metabolism, promotes catabolic processes, and regulates autophagy, oxidative stress, inflammation and cell growth. Deficient AMPK phosphorylation is a typical pathological feature of metabolic syndrome, aging-related diseases and malignant tumors.